

Product Data

D12266B



Description

Condensed milk diet. Purified match to C11024.

Used in Research

Obesity
Diabetes
Hypertension

Packaging

Product is packed in 12.5 kg box. Each box is identified with the product name, description, lot number and expiration date.

Lead Time

IN-STOCK. Ready for next day shipment.

Gamma-Irradiation

Yes. Add 10 days to delivery time.

Form

Pellet, Powder, Liquid

Shelf Life

Most diets require storage in a cool dry environment. Stored correctly they should last 3-6 months.

Control Diets

D12489B
D10001

Formula

Product #	D12266B	
	gm%	kcal%
Protein	18.5	16.8
Carbohydrate	56.7	51.4
Fat	15.6	31.8
	Total	100.0
	kcal/gm	
	4.41	
Ingredient	gm	kcal
Casein, 80 Mesh	190	760
DL-Methionine	3	12
Corn Starch	215	860
Maltodextrin 10	75	300
Sucrose	290	1160
Cellulose, BW200	30	0
Butter Fat	44.2	397.8
Corn Oil	118	1062
Mineral Mix S10001	40	0
Calcium Carbonate	5.5	0
Sodium Chloride	5.5	0
Potassium Citrate, 1 H2O	13.5	0
Vitamin Mix V10001	11	44
Choline Bitartrate	2	0
FD&C Yellow Dye #5	0	0
FD&C Red Dye #40	0.1	0
Total	1042.8	4595.8



REFERENCES

2/15/06

D12266B

1. Alberts, P. et al. Selective inhibition of 11 β -hydroxysteroid dehydrogenase type 1 decreases blood glucose concentrations in hyperglycaemic mice. *Diabetologia*. 45:1528-1532, 2002.
2. Alberts, P. et al. Selective inhibition of 11 β -hydroxysteroid dehydrogenase type 1 improves hepatic insulin sensitivity in hyperglycemic mice strains. *Endocrinology*. 144:4755-4762, 2003.
3. Turnbull, A.V., et al. Selective antagonism of the NPY Y5 receptor does not have a major effect on feeding in rats. *Diabetes*. 51:2441-2449, 2002.
4. Block, M.H., et al. Discovery and optimization of a series of carbazole ureas as NPY5 antagonists for the treatment of obesity. *J. Med. Chem.* 45:3509-3523, 2002.
5. Boustany, C.M., et al. Activation of the systemic and adipose rennin-angiotensin system in rats with diet-induced obesity and hypertension. *Am. J. Physiol. Integr. Comp. Physiol.* 287: R943-R949, 2004.
6. Chandler, P.C., et al. Change in CCK-8 response after diet-induced obesity and MC3/4-receptor blockade. *Peptides*. 25:299-306, 2004.
7. Chandler, P.C., et al. Feeding response to melanocortin agonist predicts preference for and obesity from a high-fat diet. *Physiology and Behavior*. 85:221-230, 2005.
8. Chen, A.S., et al. Role of the melanocortin-4 receptor in metabolic rate and food intake in mice. *Transgenic Research*. 9:145-154, 2000.
9. Clegg, D.J., et al. Reduced anorexic effects of insulin in obesity-prone rats fed a moderate-fat diet. *Am. J. Physiol.* 288:981-986, 2005.
10. DeRuisseau, L.R., Parsons, A.D., & Overton, J.M. Adaptive thermogenesis is intact in B6 and A/J mice studied at thermoneutrality. *Metabolism*. 53:1417-1423, 2004.
11. Bobrian, A.D., et al. Effect of salt on hypertension and oxidative stress in a rat model of diet-induced obesity. *Am. J. Physiol. Renal Physiol.* 285:F619-F628, 2003.
12. Dobrian, A.D., et al. Development of Hypertension in a rat model of diet-induced obesity. *Hypertension*. 35:1009-1015, 2000.
13. Dobrian, A.D., et al. Oxidative stress in a rat model of obesity-induced hypertension. *Hypertension*. 37:554-560, 2001.
14. Dobrian, A.D. et al. Pioglitazone prevents hypertension and reduces oxidative stress in diet-induced obesity. *Hypertension*. 43: 48-56, 2004.
15. Ghibaudi, L., et al. Fat intake affects adiposity, comorbidity factors, and energy metabolism of Sprague-Dawley rats. *Obesity Research*. 10:956-963, 2002.
16. Heinrichs, Stephen. Nonexercise muscle tension and behavioral fidgeting are positively correlated with food availability/palatability and body weight in rats. *Physiology & Behavior*. 79:199-207, 2003.
17. Larden, P.J., et al. Differential influence of peroxisome proliferators-activated receptors α and β on food intake and energy homeostasis. *Diabetes*. 52:2249-2259, 2003.
18. Lauterio, T. J., et al. Plasma growth hormone secretion is impaired in obesity-prone rats before onset of diet-induced obesity. *American Journal of Physiology*. 275:E6-E11, 1998.
19. Lauterio, T.J., Bond, J.P. * Ulman, E.A. Development and characterization of purified diet to identify obesity-susceptible and resistant rat populations. *Journal of Nutrition*. 124:2172-2178, 1994.
20. Lauterio, T.J., et al. Neuropeptide Y expression and endogenous leptin concentrations in a dietary model of obesity. *Obesity research*. 7:498-505, 1999.
21. Marsh, D.J., et al. Melanin-concentrating hormone 1 receptor-deficient mice are lean, hyperactive, and hyperphagic and have altered metabolism. *PNAS*. 99:3240-3245, 2002.
22. Shen, C.P., et al. Plasma agouti-related protein level: a possible correlation with fasted and fed states in humans and rats. *Journal of Neuroendocrinology*. 14:607-610, 2002.
23. Singh, K.A., et al. Acute insulin-induced elevations of circulating leptin and feeding inhibition in lean but not obese rats. *Am. J. Regul. Integr. Comp. Physiol.* 289:373-379, 2005.
24. Singh, D.K., et al. Phosphorylation of supernatant protein factor enhances wits ability to stimulate microsomal squalene monooxygenase. *The Journal of Biological achemistry*. 278:5646-5651, 2003.
25. Strack, A.M. et. al. Regulation of body weight and carcass composition by sibutramine in rats. *Obesity Research*. 10:173-181, 2002.
26. Turnbull, A.V. Selective antagonism of the NPY Y5 receptor does not have a major effect on feeding in rats. *Diabetes*. 51:2441-2449, 2002.
27. Warden, C. H., et al. Identification of a congenic mouse line with obesity and body length phenotypes. *DOI*. 15:460-471, 2004.
28. Zhou, D., et al. Enhanced running wheel activity of both Mch 1r- and Pmch-deficient mice. *Regulatory Peptides*. 124:53-63, 2005.